arXiv:1609.03621v1 [math.CA] 12 Sep 2016

AN EXTENSION OF NIKISHIN'S FACTORIZATION THEOREM

GEOFF DIESTEL

Abstract. A Nikishin-Maurey characterization is given for bounded subsets of weak-type Lebesgue spaces. New factorizations for linear and multilinear operators are shown to follow.

1. Introduction

Let (Ω, μ) be a non-atomic probability space. The space of scalar-valued μ-measurable functions $L_0 = L_0(d\mu)$ is equipped with the topology of convergence in measure and $f = g$ in L_0 if $f(\omega) = g(\omega)$ for μ-almost every $\omega \in \Omega$.

For $0 < p < \infty$, the Lebesgue space L_p is defined by

$$f \in L_p \Leftrightarrow \|f\|_p = \Big(\int |f|^p \, d\mu \Big)^{1/p} < \infty.$$

Of course $\|\cdot\|_p$ is a quasi-norm making L_p into a quasi-Banach space and L_p is a Banach space if $1 \leq p \leq \infty$ where L_∞ is the space of essentially bounded functions equipped with the essential supremum norm.

For $0 < p < \infty$, the weak Lebesgue space $L_{p,\infty}$ is defined by

$$f \in L_{p,\infty} \Leftrightarrow \|f\|_{p,\infty} := \sup_{t>0} t\mu(\{|f| > t\})^{1/p} < \infty.$$

The quantity $\|\cdot\|_{p,\infty}$ is a (complete) quasi-norm (see [3], Ch 1.1.1).

The intent of this article is to extend the existing Nikishin-Maurey theory. Although this theory is often presented in the context of factoring operators, the foundational results of this theory are concerned with bounded subsets of non-negative measurable functions. If $\mathcal{F}$ is such a set of measurable functions and $0 < p < \infty$, Nikishin [6] characterized the existence of a positive measurable function g such that

$$\sup_{f \in \mathcal{F}} \|f/g\|_{p,\infty} < \infty.$$

The existence of such a function is equivalent to the existence of a decreasing function $C : (0, \infty) \to (0, \infty)$ such that $\lim_{t\to\infty} C(t) = 0$ and

$$\mu(\{\sup_j |c_j f_j| > t\}) \leq C(t)$$

for all $t > 0$, finitely supported sequences (f_j) from $\mathcal{F}$ and scalars (c_j) satisfying $\sum_j |c_j|^p \leq 1$. In other words, g exists if and only if

$$\big\{ \sup_j |c_j f_j| : n \in \mathbb{N}, f_1, \cdots, f_n \in \mathcal{F}, |c_1|^p + \cdots + |c_n|^p \leq 1 \big\}$$

1991 *Mathematics Subject Classification.* Primary 46E30, Secondary 28A25.
Key words and phrases. Factorization, Type, Cotype, Banach space.

is bounded in L_0. The key step in proving g's existence from the boundedness of this set is purely constructive. A nice presentation of the proof of Nikishin's [6] characterization may be found in Proposition III.H.2 of [8]. To summarize, for $0 < \epsilon < 1$, the boundedness of the above set within L_0 implies the existence of a subset E_ϵ of Ω and a constant $C_\epsilon < \infty$ such that $\mu(E_\epsilon) \geq 1 - \epsilon$ and

$$\sup_{f \in \mathcal{F}} \|1_{E_\epsilon} f\|_{p,\infty} \leq C_\epsilon.$$

Of course the constant C_ϵ is tied to the function C from the above maximal estimate and thus it may be that $\lim_{\epsilon \to 0} C_\epsilon = \infty$. If this limit were finite there would be nothing to prove as the set $\mathcal{F}$ would be bounded in $L_{p,\infty}$. Given any decreasing null sequence (ϵ_n) such that $0 < \epsilon_n < 1$, g is constructed by selecting an unbounded increasing sequence of positive scalars (a_n) such that $g = a_1 1_{E_{\epsilon_1}} + \sum_{n>1} a_n 1_{E_{\epsilon_n} \setminus E_{\epsilon_{n-1}}}$ where the scalars (a_n) are chosen to counteract the growth of constants (C_{ϵ_n}) in order to obtain the factorization $\mathcal{F} = g(g^{-1}\mathcal{F})$ where $g^{-1}\mathcal{F}$ is bounded in $L_{p,\infty}$. Of course, any selection of scalars (a_n) defines g as a positive element of L_0. However, it is of interest to identify a more specific space for g based on the particular function $C(t)$ from the above maximal estimate. Theorem 1.1 is a formulation of Nikishin's theorem for the specific case that $\mathcal{F}$ is a subset of $L_{q,\infty}$ for some $0 < q < p$. In this case, it is natural to consider $C(t) = t^{-q}$. Much of the proof follows from a technical adaptations of the arguments used to prove Nikishin's [6] theorem as presented in [8]. However, the aforementioned construction of the function g is insufficient to obtain g from the natural weak-Lebesgue space and an extra compactness argument will be used to prove $N3 \Rightarrow N1$.

Theorem 1.1. *Let $0 < q < p < \infty$, $1/r = 1/q - 1/p$ and $\mathcal{F}$ be a subset of non-negative elements of $L_{q,\infty}$. Then the following conditions are equivalent.*

N1. There exists a constant $C < \infty$ and a positive $g \in L_{r,\infty}$ so that $\|g\|_{r,\infty} = 1$ and

$$\sup_{f \in \mathcal{F}} \|f/g\|_{p,\infty} \leq C.$$

N2. There exists a constant $C < \infty$ so that

$$\|\sup_j |c_j f_j|\|_{q,\infty} \leq C\Big(\sum_j |c_j|^p\Big)^{1/p}$$

for all finitely supported sequences (f_j) from $\mathcal{F}$ and scalars (c_j).

N3. There exists a constant $C < \infty$ such that for any $0 < \epsilon < 1$ there exists a measurable set E_ϵ such that $\mu(E_\epsilon) \geq 1 - \epsilon$ and

$$\sup_{f \in \mathcal{F}} \|1_{E_\epsilon} f\|_{p,\infty} \leq C\epsilon^{-1/r}.$$

Theorem 1.1 will be proven in Section 3. Section 2 contains applications related to the factorization of linear and multilinear operators. For more on Nikishin-Maurey theory and its applications to Banach space theory, the reader is referred to [1], [8], [9], and [10].

2. Factoring Operators

The topology of any locally bounded topological vector space X is induced by a function $\|\cdot\| : X \to [0, \infty)$ satisfying the following conditions.

(1) $\|x\| > 0$ for all $x \neq 0$.

(2) $\|cx\| = |c|\|x\|$ for all scalars c and all $x \in X$.
(3) There exists $1 \leq C < \infty$ so that $\|x+y\| \leq C(\|x\| + \|y\|)$ for all $x, y \in X$.

The function $\|\cdot\|$ is called a quasi-norm and X is a quasi-Banach space if X is complete with respect to $\|\cdot\|$. Furthermore, if $C = 1$ then X is a Banach space.

For $0 < p \leq 2$, a quasi-Banach space X has Rademacher type p if there exists a constant $T_p(X) < \infty$ such that

$$\mathbb{E}\Big\| \sum_j \epsilon_j x_j \Big\|^p \leq T_p(X)^p \sum_j \|x_j\|^p$$

for all finitely supported sequences (x_j) from X. Here (ϵ_j) is a sequence of independent Bernoulli random variables satisfying

$$P(\epsilon_j = 1) = 1/2 = P(\epsilon_j = -1).$$

For $k \geq 1$ and quasi-Banach space(s) $X_1, \cdots, X_k$, $T : X_1 \times \cdots \times X_k \to L_0$ is k-sublinear if each coordinate map is sublinear. If each coordinate map is linear then T is a k-linear operator. In [2] it was shown that for a bounded k-sublinear operator $T : X_1 \times \cdots \times X_k \to L_0$ there exists a decreasing function $C : (0, \infty) \to (0, \infty)$ such that $\lim_{t\to\infty} C(t) = 0$ and

$$\mu(\{\sup_j |T(x_{1,j}, \cdots, x_{k,j})| > t\}) \leq C(t)$$

for all finitely supported sequences $(x_{i,j})_j$ from X_i, $1 \leq i \leq k$, such that

$$\sum_j (\|x_{1,j}\| \cdots \|x_{k,j}\|)^p \leq 1$$

where $1/p = 1/p_1 + \cdots + 1/p_k$ and $0 < p_1, \cdots, p_k \leq 2$ are the respective Rademacher types of the quasi-Banach spaces $X_1, \cdots, X_k$. Thus, by Nikishin's [6] characterization there exists a positive measurable g so that

$$\|g^{-1} T(x_1, \cdots, x_k)\|_{p,\infty} \leq \|x_1\| \cdots \|x_k\|$$

for all $x_i \in X_i$, $1 \leq i \leq k$. Of course the case $k = 1$ is due to Nikishin [6]. Analogous results are also shown in [2] for operators mapping into L_q for some $0 < q < p$. These results utilize Pisier's [7] characterization for the factorization of subsets of L_q through $L_{p,\infty}$ if $0 < q < p$.

Maurey [5] characterized when a subset of L_q may be factored through L_p for $0 < q < p < \infty$. Due to the stable laws and the Kahane-Khintchine inequalities, this result only applies to linear operators defined on a space of type 2, i.e. every linear operator $T : X \to L_q$ factors through L_2 if $0 < q < 2$. There are no direct applications of Maurey's [5] results for factoring multilinear operators through L_p using Rademacher type. However, partial analogs of Maurey's [5] result are shown to hold in [2] and [4]. Suppose X_1 and X_2 have type 2 and $T : X_1 \times X_2 \to L_0$ is a bounded bilinear operator. Then not only does T factor through $L_{1,\infty}$ but the range of T is locally convex in L_0. This is shown in [2] as an application of the characterization of the Rademacher decoupling property for quasi-Banach spaces given in [4]. Thus, the T-induced linear map is continuous from the projective tensor-product $X_1 \widehat{\otimes} X_2$ into L_0. This means that T factors through a Banach space, i.e. there exists a Banach Space Z, a bilinear operator $B : X_1 \times X_2 \to Z$ and a linear operator $L : Z \to L_0$ such that $T = LB$.

As with previously existing Nikishin-Maurey theorems, Theorem 1.1 applies to linear and multilinear operators due to their homogeneity (see [2]). Suppose $k \geq 1$

and $0 < p_1, \cdots, p_k \leq 2$ are the respective Rademacher types of the quasi-Banach spaces $X_1, \cdots, X_k$. If $1/p = 1/p_1 + \cdots + 1/p_k$ and $T : X_1 \times \cdots \times X_k \to L_{q,\infty}$ is a bounded k-linear operator and $0 < q < p$ then the Kahane-Khintchine inequalities imply that there exists a constant $C = C(q, p_1, \cdots, p_k, T) < \infty$ such that

$$\Big\| \Big(\sum_{j_1, \cdots, j_k} |T(x_{1,j_1}, \cdots, x_{k,j_k})|^2 \Big)^{1/2} \Big\|_{q,\infty} \leq C \prod_{i=1}^{k} \Big(\sum_{j_i} \|x_{i,j_i}\|^{p_i} \Big)^{1/p_i}$$

for all finitely supported sequences $(x_{i,j})_j$ from X_i, $1 \leq i \leq k$. Of course this estimate implies that

$$\| \sup_j |T(x_{1,j}, \cdots, x_{k,j})| \|_{q,\infty} \leq C \prod_{i=1}^{k} \Big(\sum_{j} \|x_{i,j}\|^{p_i} \Big)^{1/p_i}$$

for all finitely supported sequences $(x_{i,j})_j$ from X_i, $1 \leq i \leq k$. As illustrated in [4] and [2], the homogeneity of T and of the identity $1/p = 1/p_1 + \cdots + 1/p_k$ implies

$$\| \sup_j |T(x_{1,j}, \cdots, x_{k,j})| \|_{q,\infty} \leq C \Big(\sum_{j} (\|x_{1,j}\| \cdots \|x_{k,j}\|)^p \Big)^{1/p}$$

for all finitely supported sequences $(x_{i,j})_j$ from X_i, $1 \leq i \leq k$. An immediate consequence of this estimate and Theorem 1.1 is the following.

Theorem 2.1. *Let $k \geq 1$, $0 < p_1, \cdots, p_k \leq 2$, $1/p = 1/p_1 + \cdots + 1/p_k < 1/q < \infty$, $1/r = 1/q - 1/p$ and suppose $T : X_1 \times \cdots \times X_k \to L_{q,\infty}$ is a continuous k-linear operator where $p_1, \cdots, p_k$ are the respective Rademacher types of the quasi-Banach spaces $X_1, \cdots, X_k$. Then there exists a positive $g \in L_{r,\infty}$ such that*

$$\|g^{-1} T(x_1, \cdots, x_k)\|_{p,\infty} \leq \|x_1\| \cdots \|x_k\|$$

for all $x_i \in X_i$, $1 \leq i \leq k$.

Theorem 2.1 identifies new factorizations not established by the characterizations of Nikishin [6], Maurey [5], or Pisier [7] due the the fact that the operator takes values in $L_{q,\infty}$ and that the factorization $T = g(g^{-1}T)$ is defined by an element of g of $L_{r,\infty}$ where $1/r = 1/q - 1/p$.

3. Proof of Theorem 1.1

All Nikishin-Maurey characterizations like Theorem 1.1 may be reduced to any particular value of p by considering

$$\mathcal{F}^t = \{ f^t : f \in \mathcal{F} \}$$

for any $t > 0$. Since $1/r = 1/q - 1/p$, it follows that $1/(r/t) = 1/(q/t) + 1/(p/t)$ and all the conditions $N1 - N3$ translate into equivalent conditions about $\mathcal{F}^t$. Picking t so that p/t has a particular value and proving the equivalence of these new conditions for $\mathcal{F}^t$ implies the general equivalence of $N1 - N3$.

The following proof of $N1 \Rightarrow N2 \Rightarrow N3$ follows from a technical reworking of the arguments used to prove the analogous implications of Nikishin's [6] theorem as presented in [8]. However, the aforementioned constructive arguments used to prove the final implication of Nikshin's [6] theorem are insufficient to construct the function g from $N1$ because of the added requirement that $g \in L_{r,\infty}$ where

$1/r = 1/q - 1/p$. However, assuming $N3$ and using the aforementioned reduction with the assumption that $r > 2$, one may construct a sequence (g_n) such that

$$\sup_n \sup_{f\in\mathcal{F}} \|f/g_n\|_{p,\infty} < \infty$$

and

$$\sup_n \|g_n\|_{r_n,\infty} < \infty$$

where $2 < r_n < r$ and (r_n) is increasing to r. Thus, (g_n) is bounded in L_2 and the Banach-Saks Theorem guarantees that (g_n) has a subsequence with convergent Cesàro means. The above two conditions will be used to show that the limit g of these Cesàro means will satisfy $N1$ and this will complete the proof by establishing $N3 \Rightarrow N1$.

Proof. Assuming $N1$, let (f_j) be a finitely supported sequence from $\mathcal{F}$, (c_j) be scalars, and define $F = \sup_j |c_j f_j|$. Then $1/r = 1/q - 1/p$ implies

$$\|F\|_{q,\infty} \le \|F/g\|_{p,\infty}\|g\|_{r,\infty}.$$

Since (f_j) is a finitely supported sequence from $\mathcal{F}$, there is a partition (E_j) of Ω so that

$$F = |c_k f_k|$$

almost everywhere on E_k. The disjointness of the partition implies

$$\mu(\{F/g > t\}) \le \sum_k \mu(\{|c_k f_k|/g > t\}).$$

Hence, $N1$ implies

$$\mu(\{\sup_j |c_j f_j|/g > t\}) \le \sum_k \mu(\{|c_k f_k| > t\}) \le Ct^{-p} \sum_k |c_k|^p$$

and $N2$ follows with same constant as in $N1$ because $\|g\|_{r,\infty} = 1$.

Now assume $N2$ holds. Without loss of generality, assume $C = 1$ by normalizing $\mathcal{F}$. Fix $0 < \epsilon < 1$ and consider a measurable set B to be ϵ-bad if there exists $f \in \mathcal{F}$ so that

$$\mu(B)|f(\omega)|^p > \epsilon^{-p/q}$$

for all $\omega \in B$.

If there are no ϵ-bad sets for some $0 < \epsilon < 1$ then for every $f \in \mathcal{F}$ and every $t > 0$, there exists $\omega \in \{|f| > t\}$ so that

$$\mu(\{|f| > t\})t^p < \mu(\{|f| > t\})|f(\omega)|^p \le \epsilon^{-p/q}.$$

In this case $\mathcal{F}$ is bounded in $L_{p,\infty}$ and $N3$ follows in a trivial manner.

Suppose there are ϵ-bad sets for every $0 < \epsilon < 1$. Fix ϵ and suppose (B_j) is a maximal family of disjoint ϵ-bad sets. Thus, for each j there exists $f_j \in \mathcal{F}$ so that

$$\mu(B_j)|f_j(\omega)|^p > \epsilon^{-p/q}$$

for all $\omega \in B_j$. For each j let $c_j = \mu(B_j)^{1/p}$. Notice that

$$\sup_j |c_j f_j|^p > \epsilon^{-p/q}$$

everywhere on $B = \cup_j B_j$. Thus, for any $n \in \mathbb{N}$, $N2$ implies that

$$\mu(\cup_{j=1}^n B_j) \le \mu(\{\sup_{j\le n} |c_j f_j| > \epsilon^{-1/q}\}) \le \epsilon\Big(\sum_{j=1}^n |c_j|^p\Big)^{q/p} \le \epsilon\mu(B)^{q/p}.$$

Letting n tend to infinity implies that

$$\mu(B)^{1-q/p} \leq \epsilon.$$

Therefore,

$$\mu(B) \leq \epsilon^{p/(p-q)}.$$

So, if $E_{\epsilon^{p/(p-q)}} = \Omega \setminus B$ then $\mu(E_{\epsilon^{p/(p-q)}}) \geq 1 - \epsilon^{p/(p-q)}$ and $E_{\epsilon^{p/(p-q)}}$ is not ϵ-bad. Therefore,

$$\|1_{E_{\epsilon^{p/(p-q)}}} f\|_{p,\infty}^p \leq \epsilon^{-p/q}.$$

This holds for all $0 < \epsilon < 1$. By making the substitution $\delta = \epsilon^{p/(p-q)}$ then for every $0 < \delta < 1$, the condition $1/r = 1/q - 1/p$ implies that there exists E_δ such that $\mu(E_\delta) \geq 1 - \delta$ and

$$\|1_{E_\delta} f\|_{p,\infty}^p \leq \delta^{-(p-q)/q} = \delta^{-p/r}.$$

By undoing the assumed normalization of $N2$, it follows that $N3$ holds with the same constant as in $N2$.

Assume $N3$ with $C = 1$. Moreover, the fact that $\mathcal{F}$ is a set of positive measurable functions, the homogeneity of the desired estimates on the indices q, p, and r due to the identity $1/r = 1/q - 1/p$ allows for the assumption that $r > 2$.

Fix n and let $(\epsilon_{n,m})_m$ be the sequence defined by $\epsilon_{n,m} = \big(1/(2m^n)\big)_m$. For each m, let $E_{n,m}$ be the set $E_{\epsilon_{n,m}}$ from condition $N3$ for $\epsilon = \epsilon_{n,m}$. The decreasing nature of the sequence $(\epsilon_{n,m})_m$ implies we may assume (with no loss of generality) that

$$E_{n,m} \subset E_{n,m+1}$$

for all n and m. Moreover, since μ is non-atomic, we may assume $\mu(E_{n,m}) = 1 - 1/(2m^n)$ by choosing $E_{n,m}$ as a subset of $E_{\epsilon_{n,m}}$ if necessary. Thus, for every n and each $m \geq 1$,

$$\mu(E_{n,m+1} \setminus E_{n,m}) = \tfrac{1}{2m^n} - \tfrac{1}{2(m+1)^n} = \tfrac{(m+1)^n - m^n}{2m^n(m+1)^n} \leq \tfrac{n}{2m^{n+1}}.$$

Let $D_{n,1} = E_{n,1}$ and $D_{n,m} = E_{n,m+1} \setminus E_{n,m}$ for all $m > 1$.

Suppose the above construction has been made for all n. For each n, define a measurable function g_n by

$$g_n = \sum_m 1_{D_{n,m}} m^{(n+k)/r}$$

where k is chosen so that $pk/r > 1$. Clearly, $g_n \geq 1$ on Ω for all n.

Let $r_n = \frac{rn}{n+k}$. Then (r_n) increases to r. Furthermore, it is easy to see that there is a constant $C < \infty$ which is independent of n so that

$$\mu(\{g_n > t\}) = \sum_{m > t^{r/(n+k)}} \mu(D_{n,m}) \leq \sum_{m > t^{r/(n+k)}} \tfrac{n}{2m^{n+1}} \leq Ct^{-rn/(n+k)}.$$

Thus, $\sup_n \|g_n\|_{r_n,\infty} < \infty$. Moreover, if $f \in \mathcal{F}$ then

$$\mu(\{|f|/g_n > t\}) = \sum_m \mu(D_{n,m} \cap \{|f| > tm^{(n+k)/r}\}).$$

Since $D_{n,m} \subset E_{\epsilon_{n,m}}$, $N3$ and our choice of k imply that there is a constant $C = C(p,q,r) < \infty$ which is independent of n so that

$$\mu(\{|f|/g_n > t\}) \leq t^{-p} \sum_m 2^{p/r} m^{np/r} m^{-p(n+k)/r} = t^{-p} 2^{p/r} \sum_m m^{-pk/r} = Ct^{-p}.$$

Since $r > 2$ and (r_n) is increasing to r, we may restrict our attention to all n such that $2 < r_n < r$. Without loss of generality, we may assume that $r_1 > 2$. Since μ is a probability measure

$$\sup_n \|g_n\|_2 \leq C(r_1) \sup_n \|g_n\|_{r_n,\infty} < \infty.$$

By the Banach-Saks Theorem for L_2, there exists a subsequence $(g_{n_k})_k$ and $g \in L_2$ so that

$$g = \lim_N G_N \quad (in\ L_2)$$

where

$$G_N = N^{-1} \sum_{j=1}^{N} g_{n_j}.$$

For any $m \geq 1$ define

$$G_{N,m} = N^{-1} \sum_{j=m}^{N+m-1} g_{n_j}.$$

By a simple application of the triangle inequality, it is easy to see that g is the L_2 limit of $(G_{N,m})_N$ for any m. By Fatou's lemma,

$$\mu(\{g > t\}) \leq \liminf_N \mu(\{G_{N,m} > t\})$$

for any m. However, since (r_n) is increasing and μ is a probability measure, notice that $G_{N,m}$ is an average of functions whose $L_{r_m,\infty}$ norms are uniformly bounded in m by a constant independent of N. So, there exists a constant $C(p,q) < \infty$ so that

$$\sup_m \|g\|_{r_m,\infty} < C(p,q)$$

Since μ is a probability measure and (r_m) increases to r, it follows that $g \in L_{r,\infty}$ with $\|g\|_{r,\infty} \leq C(p,q)$.

Fix $f \in \mathcal{F}$. Again, by applying Fatou's lemma,

$$\mu(\{|f|/g > t\}) \leq \liminf_N \mu(\{|f|/G_N > t\}).$$

By the arithmetic-geometric mean inequality

$$G_N \geq \Big(\prod_{j=1}^{N} g_{n_j} \Big)^{1/N}.$$

Therefore, by Hölder's inequality for weak-type spaces,

$$\mu(\{|f|/G_N > t\}) \leq t^{-p} \||f|/G_N\|_{p,\infty}^p \leq t^{-p} \prod_{i=1}^{N} \||f|^{1/N}/g_{n_i}^{1/N}\|_{Np,\infty}^p = t^{-p} \prod_{i=1}^{N} \|f/g_{n_i}\|_{p,\infty}^{p/N}.$$

By normalizing g in $L_{r,\infty}$, the condition $\sup_n \sup_{f\in\mathcal{F}} \|f/g_n\|_{p,\infty} < \infty$ implies there is a constant $C < \infty$ such that

$$\sup_{f\in\mathcal{F}} \|f/g\|_{p,\infty} \leq C$$

and $N1$ follows.

□

References

[1] F. Albiac and N. Kalton, *Topics in Banach spaces*,Springer, New York (2006).

[2] G. Diestel, *Factoring multisublinear maps*, J. Functional Analysis **266** issue 4 (2014), 1928–1947.

[3] L. Grafakos, *Classical Fourier Analysis,*Second Edition, Graduate Texts in Math., no 249, Springer, New York, 2008.

[4] N.J. Kalton, *Rademacher series and decoupling*,New York J. of Math. **11**(2005), 563–595.

[5] B. Maurey, *Théorèms de factorisation pour les opérateurs linéaires à valeurs dans les epaces* L_p,Société Mathemátique de France, Paris, 1974, With an English summary; Astérisque, No. 11 .(French)

[6] E.M. Nikishin, *Resonance theorems and superlinear operators*, Uspekhi Mat. Nauk 25 No. 6 (1970), 129–191.(Russian)

[7] G. Pisier, *Factorization of Operators Through* $L_{p,\infty}$ *or* $L_{p,1}$ *and Non-Commutative Generalizations,*Math. Ann. **276** (1986), 105–136.

[8] P. Wojtaszczyk, *Banach spaces for analysts.*Cambridge Studies in Advanced Mathematics, vol. 25, Cambridge University Press, Cambridge, 1991.

[9] *Handbook of the Geometry of Banach Spaces, Vol. 1.* Edited by W.B. Johnson and J. Lindenstrauss. North-Holland Publishing Co., Amsterdam, 2001.

[10] *Handbook of the Geometry of Banach Spaces, Vol. 2.* Edited by W.B. Johnson and J. Lindenstrauss. North-Holland Publishing Co., Amsterdam, 2001.

Texas A&M University-Central Texas, 1001 Leadership Place, Killeen TX 76549, USA

E-mail address: gdiestel@tamuct.edu